

Human Papillomavirus (HPV) and Cytology Tests Requisition – For Cervical Screening

- Eligibility Criteria: People with a cervix age 25 and older who have ever been sexually active and have a valid OHIP number.
- Ontario Cervical Screening Program's cervical screening recommendations and cessation criteria can be found at ontario-number-12
- Immunocompromised populations include people who are living with HIV/AIDS (regardless of CD4 cell count), congenital (primary) immunodeficiency, systemic lupus erythematosus (regardless of whether they are receiving immunosuppressant treatment), renal failure and require dialysis, transplant recipients (solid organ or allogeneic stem cell transplants) or people requiring treatment (either continuously or at frequent intervals) with medications that cause immune suppression for 3 years or more.

Lab Use Only

• Referral to a specialist is required for any visible cervical abnormalities	es.
Requester Information Requester type (check ONE): Physician Midwife Nurse practitioner	Patient Identification (Enter information as indicated on OHIP card. Can be replaced by a sticker.)
CPSO or CNO number:	Last name:
Practitioner billing number:	Middle name: (optional)
Last name:	First name:
Middle name: (optional)	Date of birth: Sex: Male Female
First name:	OHIP number: OHIP version:
Address:	Patient Contact (Mailing address for result letters and other
Fax: () Phone: ()	correspondence. Verify with patient.)
Copy to: Primary care provider	Building / Street name:
Last name:	Apt./Unit number: City:
First name:	Province: Postal Code:
Address: (optional)	Phone: () Extension: (optional)
Fax: () Phone: ()	Type: Home Work Cell
Testing Indication for Cervical Screening (check ONE):	Specimen
A. HPV test (includes reflex cytology if HPV-positive)	Site: Cervical/endocervical Vaginal Double cervix
Average risk screening: every 5 years	Special considerations for cytology interpretation:
Immunocompromised screening: every 3 years	Intrauterine device (IUD) Postpartum
HPV-positive (other high-risk types) with normal or low-grade (NILM/ASCUS/LSIL) cytology: 2-year follow-up (moderate risk)	Menopausal hormone Pregnancy therapy (MHT) Subtotal hysterectomy
More frequent screening post-colposcopy: 2-year follow-up (moderate risk)	Post-menopausal Transition-related hormone therapy
People with histologic evidence of dysplasia in the cervix at the time of hysterectomy and people with a history of early cervical cancer: 1-time post-hysterectomy vaginal vault testing	Specimen collection date: (yyyy/mm/dd)
	Last menstrual period (first day): (yyyy/mm/dd)
B. Cytology test only	Clinical information
Repeat after a previous HPV-positive (other high-risk types) with unsatisfactory cytology result	
Requester Verification	Date:
Requester signature:	(yyyy/mm/dd)